

HEALTH SCIENCES IMMUNIZATION PROGRAM (HSIP) REQUIREMENTS CHECKLIST

This document provides a summary of the Health Sciences Immunization Program (HSIP) requirements. We recommend taking this to your provider when completing your requirements. Detailed instructions are on the [HSIP webpage](#) and will be available in your Med+Proctor account.

DOCUMENT SUBMISSION PROCESS

Students submit their documentation to a web-based system called Med+Proctor. Once accepted into your HSIP-participating program, your program will send you instructions on how to create your Med+Proctor account and specific due dates for meeting your HSIP requirements. **Please wait to create your Med+Proctor account until your program has given you the instructions.**

Submit records into Med+Proctor as you complete an HSIP requirement. This way, the tags in your account are up to date with what you need to complete next. If you are in process of completing a vaccine series, submit documentation as you complete each dose.

CHILDHOOD IMMUNIZATIONS

The Childhood Immunization requirement is an attestation confirming receipt of a primary childhood tetanus, diphtheria, pertussis (DTaP/DTP/DT/Td) AND Polio (IPV/OPV) vaccine series. **Documentation is not required.** You will attest "Yes" if you have received BOTH childhood vaccine series. You will attest "No" if you have not received either or have received only one of the series. Contact HSIP at myshots@uw.edu if you answer "No."

MMR: MEASLES (RUBEOLA), MUMPS AND RUBELLA

Two doses of the MMR vaccine are required (other vaccine types are not accepted) **OR** positive measles, mumps, and rubella IgG antibody titers (Note: IgM titer is not acceptable). You can complete the requirement through the following:

- ☐ Two doses of MMR vaccine dated January 1971 or later and given on or after 12 months of age. Doses must be at least 28 days apart. Live vaccines may be given on the same day otherwise they must be spaced at least 28 days apart (e.g., MMR and Varicella); **OR**

- ☐ Lab reports showing positive IgG antibody titer results for measles (rubeola), mumps, and rubella.

VARICELLA

Two doses of the Varicella vaccine are required, **OR** positive Varicella IgG antibody titer (Note: IgM titer is not acceptable). **History of disease is not accepted.** You can complete the requirement through the following:

- ☐ Two doses of Varicella vaccine dated March 1995 or later and given on or after 12 months of age. Doses must be at least 28 days apart. Live vaccines may be given on the same day otherwise they must be spaced at least 28 days apart; **OR**
- ☐ Lab report showing positive varicella IgG antibody titer result.

TETANUS-DIPHTHERIA-PERTUSSIS

One adult dose of **Tdap** completed within the last 10 years is required. **Titers are not accepted in lieu of Td/Tdap vaccine.** You can complete the requirement through the following:

- ☐ Documentation of an adult Tdap vaccine within the past 10 years; **OR**
- ☐ Documentation of an adult Td within the past 10 years **AND** a Tdap vaccine after 6/1/2005.

MENINGITIS ACWY VACCINE (GENERAL UW IMMUNIZATION REQUIREMENT)

If you are an HSIP student aged 16 through 21, then you must satisfy the [General UW Immunization Meningitis ACWY \(meningococcal conjugate\) requirement](#). You are required to submit documentation of a Meningitis ACWY vaccine taken on or after your 16th birthday. This requirement is managed by the UW Immunity Verification Program (IVP), not HSIP.

ANNUAL INFLUENZA VACCINE

An annual fall seasonal influenza vaccine is required **after** August of your entry year and each year thereafter. An Influenza vaccine requirement tag will populate in your Med+Proctor account every August. Please refer to the [HSIP webpage](#) for your requirement's due date.

TUBERCULOSIS (TB) SCREENING (NOT A VACCINE)

An initial (entry year) screening of an **IGRA blood test (QuantiFERON-TB Gold or T-Spot lab test)** is **strongly preferred and should be completed but those who have history of the BCG vaccine**. Refer to the [HSIP webpage](#) for an acceptable screening time frame and due date.

TWO separate TB skin tests (PPD) are also acceptable. The second PPD must be placed between 1-3 weeks after the first is read. PPD documentation must be from a verifiable medical source and include full name, dates placed, dates read, and results in millimeter (mm).

Previous or current positive TB screening requires a copy of the positive TB screening (positive IGRA full lab report or positive PPD skin test record), a chest x-ray report completed **AFTER** the positive TB screening, and completion of the HSIP TB Symptom Survey form.

DOCUMENTATION REQUIREMENTS

- All records must be from a medically verifiable source and include full student name.
- Vaccine records must clearly indicate vaccine type, and exact dates (MM/DD/YYYY) for each dose. Pharmacy printouts, provider/clinic-verified childhood records/cards, chart notes, and state immunization information/registry records are all acceptable proof of immunization.
- Lab results require a full lab report and must include the date of collection. A healthcare employee immunization summary is accepted for MMR & Varicella titers ONLY. A numeric reference range must be included for the Hepatitis B titer.

HEPATITIS B

BOTH documentation of a fully completed Hepatitis B vaccine series spaced appropriately per standards set by the Centers for Disease Control and Prevention (CDC) **AND a POST-vaccination positive Quantitative Hepatitis B Surface Antibody Titer** is required.

- ❑ **Step 1:** Documented Hepatitis B vaccine series (Recombivax HB, Energix-B, Twinrix, or other three dose vaccine series) **OR** Heplisav-B 2-dose vaccine series. Students unable to locate past Hepatitis B vaccine records should complete a 2-dose Heplisav-B vaccine series. If completing a series, upload documentation of each vaccine dose into Med+Proctor as it is received.
- ❑ **Step 2:** A positive Quantitative Hepatitis B Surface Antibody Titer (HBsAb-QN/CONC/Index titer) drawn at least 1 month **after** completion of a documented vaccine series. A full lab report is required (see [Documentation Requirements section](#)).

Negative/Non-Reactive Hepatitis B Surface Antibody Titer – indicates you do not have proven immunity to Hepatitis B. Proceed with getting a single additional vaccine dose (Heplisav-B highly recommended) followed by a repeat Quantitative Hepatitis B Surface Antibody Titer 1 month after the additional vaccine dose. If the repeat titer is still negative, complete the remaining vaccine doses in the 2nd series AND another titer 1 month after the final dose.

Hepatitis B non-responders – Individuals with a negative Quantitative Hepatitis B Surface Antibody titer following TWO documented Hepatitis B vaccine series are considered non-responders to the vaccine. Proceed with getting a Hepatitis B Antigen Titer and contact HSIP at myshots@uw.edu.

Hepatitis B Infection (acute, chronic, or resolved) – Contact HSIP at myshots@uw.edu.